



WHAT IS WHOOPING COUGH? (PERTUSSIS)

Whooping cough (or pertussis) is a **highly** contagious respiratory infection caused by the bacterium *Bordetella pertussis*

- It can affect babies, children, adolescents and adults.
- For adolescents and adults the infection may only cause an irritating, persistent cough. However, whooping cough can be life threatening for babies and young children, particularly those not fully protected by vaccination. During coughing attacks, a baby or child's breathing can be obstructed and they may become blue or stop breathing.

Symptoms

- may start like a cold, with a runny nose, sneezing and tiredness, and then the characteristic cough develops.
- coughing bouts can be very severe and frightening, and may end with a crowing noise (the whoop). This occurs as air is drawn back into the chest, and can be followed by vomiting or gagging.
- Bouts of coughing may continue for many weeks even after treatment.

Transmission

Whooping cough bacteria are highly infectious and are spread to other people by an infected person coughing and sneezing. The infection can also be passed on through direct contact with infected secretions from the mouth or nose. The time between exposure to the bacteria and getting sick is usually seven to ten days, but can be up to three weeks.

A person is most infectious in the early stages of their illness. Unless treated with antibiotics, a person is regarded as infectious for three weeks after the cough began.

Treatment

- a full course of antibiotics which reduces the time a person is infectious to others.
- Antibiotics need to be given within 21 days of the start of general symptoms or within 14 days of the start of coughing. Antibiotics may reduce symptoms if given early.

Some people who have had close contact with an infected person may need to take

antibiotics to prevent infection. This includes people at high risk of serious complications (eg. children aged less than one year who are not fully vaccinated, pregnant women and the elderly) and others who live or work with people at risk.

Control

- Stay away from work, school, preschool and child-care until they have had at least 5 days of their course of antibiotics, or until 21 days after the cough began.
- Children with less than three doses of whooping cough vaccine are close contacts of an infected person, they should stay away from preschool and childcare until they have taken **a full course of antibiotics, or for 14 days after their last exposure** to the infected person.

Prevention

Vaccination is the most effective way to prevent whooping cough. The vaccine is recommended and available free for small children and those in Year 10 (booster)

A booster dose is also recommended for any adults who haven't had a previous booster, particularly:

- parents planning a pregnancy, or as soon as the baby is born
- anyone working with or caring for babies and young children, especially healthcare workers, childcare workers and grandparents.

Help and Assistance

For further information please contact your local doctor, community health centre or nearest public health unit or hospital.



For further information and advice contact the Workers Health Centre



WORKERS HEALTH CENTRE

People | Recovery | Work

Industrial Health and Research Foundation - ABN 50 804 045 194

Address Ground Floor, Suite 1, 20 – 24 Wentworth Street Parramatta NSW 2150

Tel (02) 9749 7666 **Fax** (02) 9897 2488 **Email** admin@workershealth.com.au **Website** www.workershealth.com.au

WHC acknowledges support in the production of these Factsheets, from the Western Sydney Local Health District (WSLHD), a Division of the NSW Ministry of Health.

Facts042© Workers Health Centre 2016